

Platonic polyhedra tune the 3-sphere: Harmonic analysis on simplices.

Peter Kramer, Institut fuer Theoretische Physik,
U Tuebingen, Germany.

January 5, 2009

Abstract.

A spherical topological manifold of dimension $n - 1$ forms a prototile on its cover, the $(n-1)$ -sphere. The tiling is generated by the fixpoint-free action of the group of deck transformations. By a general theorem, this group is isomorphic to the first homotopy group. A basis for the harmonic analysis on the $(n-1)$ -sphere is given by the spherical harmonics which transform according to irreducible representations of the orthogonal group. Multiplicity and selection rules appear in the form of reduction of group representations. The deck transformations form a subgroup, and so the representations of the orthogonal group can be reduced to those of this subgroup. Upon reducing to the identity representation of the subgroup, the reduced subset of spherical harmonics becomes periodic on the tiling and tunes the harmonic analysis on the $(n-1)$ -sphere to the manifold. A particular class of spherical 3-manifolds arises from the Platonic polyhedra. The harmonic analysis on the Poincare dodecahedral 3-manifold was analyzed along these lines. For comparison we construct here the harmonic analysis on simplicial spherical manifolds of dimension $n = 1, 2, 3$. Harmonic analysis can be applied to the cosmic microwave background observed in astrophysics. Selection rules found in this analysis can detect the multiple connectivity of spherical 3-manifolds on the space part of cosmic space-time.

1 Introduction.

Viewed on its universal cover $S^{(n-1)}$, a spherical topological manifold $\mathcal{M}$ is the prototile of a tiling generated by the corresponding deck transformations. By a general theorem in topology given by Seifert and Threlfall [11] pp. 195-8, the group $deck(\mathcal{M})$

of deck transformations is isomorphic to the first homotopy group $\pi_1(\mathcal{M})$ and hence is a topological invariant.

A particular class of spherical 3-manifolds arises from the five Platonic polyhedra. Everitt [2] discusses their topology and gives a graphical algorithm for their homotopy groups. In Table 1.1 we list the five polyhedra and from [2] give the known order of their homotopy group.

Table 1.1. The five 3-manifolds arising from the Platonic polyhedra, the orders $|\pi_1(\mathcal{M})|$ of their first homotopy groups, and the volume fraction $frac(\mathcal{M}) = |\pi_1(\mathcal{M})|^{-1}$ of the prototile w.r.t the volume of the 3-sphere.

$\mathcal{M} :$	tetrahedron	cube	octahedron	icosahedron	dodecahedron	
$ \pi_1(\mathcal{M}) :$	5	8	8		120	(1)
$frac(\mathcal{M}) :$	0.2	0.125	0.125		0.0083..	

Since the groups $deck(\mathcal{M})$ and $\pi_1(\mathcal{M})$ are isomorphic and $deck(\mathcal{M})$ acts fixpoint-free on S^3 , the volume fraction of $\mathcal{M}$ taken as prototile on S^3 is equal to $|\pi_1(\mathcal{M})|^{-1}$. We see that the tetrahedron and the dodecahedron display extremal values of this fraction.

For general notions of topology we refer to [11], [12]. The harmonic analysis on these manifolds can be started from $S^{(n-1)}$. There its basis is the complete, orthonormal set $\langle Y^\lambda \rangle$ of spherical harmonics, the square integrable eigenmodes of $S^{(n-1)}$. To pass to a 3-manifold $\mathcal{M}$ universally covered by $S^{(n-1)}$, consider the maximal subset $\langle Y^{\lambda_0} \rangle$ of this basis periodic with respect to deck transformations. Due to the periodicity, it can be restricted to the prototile $\mathcal{M}$ and forms its eigenmodes. These periodic eigenmodes tune the sphere $S^{(n-1)}$ to the topology of $\mathcal{M}$.

To analyze in detail the periodic eigenmodes we turn to groups and their representations. Under the group $O(n, R)$ of isometries of $S^{(n-1)}$, the spherical harmonics Y^λ transform according to a set D^λ of irreducible representations. The periodic subset Y^{λ_0} transforms according to the identity representation D^0 of the group $deck(\mathcal{M})$ of deck transformations. In terms of the group/subgroup pair $O(n, R) > deck(\mathcal{M})$, we require the reduction of the irreducible representations $D^\lambda > D^0$. Clearly not all representations D^λ will reduce to the representation D^0 . The non-occurrence provides a selection rule on eigenmodes of $S^{(n-1)}$ which is another mark for the topology of $\mathcal{M}$.

Motivated by physics, harmonic analysis on topological 3-manifolds has been invoked in cosmological models of the space part of space-time [8], [9]. A direct experimental access to the topology from the autocorrelation of the cosmic matter distribution is difficult. As an alternative, the rich data from fluctuations of the Cosmic Microwave Background (CMB) radiation were examined by harmonic analysis. It is hoped to find in this way the characteristic selection rules and tuning for a specific non-trivial topology, distinct from the standard simply-connected one. Of the Platonic

3-manifolds, the Poincare dodecahedral manifold of minimal volume fraction and its eigenmodes have found particular attention. Representation theory was applied to the harmonic analysis on Poincare's dodecahedral 3-manifold in [6]. A comparative study of the harmonic analysis, tuned to different topological 3-manifolds, can provide clues for future applications.

To initiate such a comparative study, we turn here to simplicial manifolds on $S^{(n-1)}$, $n-1 = 1, 2, 3$. For illustration of the group and representation theory we start in sections 2 and 3 with the cases $n-1 = 1, 2$. Section 4 deals with the Platonic tetrahedral 3-manifold.

A regular simplex with $(n+1)$ vertices can be centrally projected to $S^{(n-1)}$ to yield a tiling into $(n+1)$ spherical simplices. The simplicial tiling can be generated by the fixpoint-free action of the cyclic group C_{n+1} acting as $\text{deck}(\mathcal{S}_0(n-1))$ on a prototile. In Fig. 1 we illustrate symbolically the tilings and simplicial manifolds for $n-1 = 1, 2, 3$.

We require the identity representation D^0 of C_{n+1} in the reduction of representations of the groups

$$O(n, R) > C_{n+1}. \quad (2)$$

Working with this pair of groups will display the difference in the topology between $S^{(n-1)}$ and $\mathcal{S}_0(n-1)$ as part of the harmonic analysis.




Fig.1 Simplicial topological manifolds on spheres. Regular simplices with $(n+1)$ vertices are inscribed into spheres $S^{(n-1)}$, $n = 2, 3, 4$, here represented symbolically by a circle. Central projection yields a simplicial tiling of $S^{(n-1)}$. Projection of the subsimplex $\mathcal{M} = \mathcal{S}_0(n-1)$, heavy lines, obtained by dropping the vertex labelled $(n+1)$, represents a simplicial topological $(n-1)$ -manifold as a prototile on its universal cover $S^{(n-1)}$. The group $\text{deck}(\mathcal{S}_0(n-1))$ is the cyclic group C_{n+1} . It acts fixpoint-free on $S^{(n-1)}$ and produces the simplicial tiling. By a general theorem it is isomorphic to the first homotopy group $\pi_1(\mathcal{S}_0(n-1))$ which is a topological invariant. Restricting the basis of the harmonic analysis from $S^{(n-1)}$ to its subset periodic under C_{n+1} we construct the harmonic analysis on $\mathcal{S}_0(n-1)$.

Now we note that the original n -simplex, of which $\mathcal{S}_0(n-1)$ forms a spherical face, has the symmetry group $S(n+1)$, the symmetric group on $(n+1)$ variables. The representation theory of $S(n+1)$ is well known [10] pp. 38-9, [3] pp. 214-31, and we shall make full use of it.

$S(n+1)$ is a Coxeter group generated by reflections, see [4]. It has the Coxeter diagram with n nodes

$$S(n+1) = \circ - \circ - \dots - \circ \quad (3)$$

Each node of this diagram, [4] pp. 31-3, describes the generator of a Weyl reflection W_a determined by a Weyl or reflection vector a in the Euclidean space E^n embedding $S^{(n-1)}$, with W_a the involutive map

$$x \in E^n : (W_a \times x) \rightarrow x - 2 \frac{\langle x, a \rangle}{\langle a, a \rangle} a. \quad (4)$$

Moreover the diagram eq. 3 implies definite relations and scalar products between the various Weyl reflection vectors. In section 4.3 we shall take $S(5)$ as a Coxeter group and describe its Weyl reflection vectors.

Since C_{n+1} is generated by the cyclic permutation $(1, 2, \dots, n+1) \in S(n+1)$, we can refine the group/subgroup pair eq. 2 as

$$O(n, R) > S(n+1) > C_{n+1}. \quad (5)$$

The group $S(n+1)$ here is taken as a subgroup of $O(n, R)$ via the orthogonal irreducible representation $D^{[n1]}$ of partition $f = [n1]$ since (i) the Weyl reflections of the defining representation for the Coxeter group eq. 3 are orthogonal transformations, (ii) the irreducible orthogonal representations of $S(n+1)$ are characterized by Young diagrams f . The orthogonal irreducible representation $D^f, f = [n1]$ has dimension n and is equivalent to the defining representation of the Coxeter group.

The cyclic subgroup $C_{n+1} < S(n+1)$ in eq. 5 is generated by the product of all the Weyl reflection generators of the Coxeter group. This product is termed the Coxeter element, see [4] pp. 74, 174.

An advantage in using the scheme eq. 5 is the fact that the subgroup $S(n) < S(n+1)$ permuting the vertices $1, 2, \dots, n$ naturally appears as a symmetry subgroup acting on the spherical simplex $\mathcal{S}_0(n)$. Borrowing the terminology from space groups in E^n , we call $S(n)$ the point group of $\mathcal{S}_0(n)$.

The reduction of irreducible representations for $O(n, R) > S(n+1)$ was studied in [5] in full detail for $n = 2, 3$. This allows to work out the harmonic analysis for these cases. We apply the representation theory of groups, following [5], in the following sections to the harmonic analysis on the simplicial manifolds for $n-1 = 1, 2, 3$. By a first step in the reduction of representations according to eq. 5 we find multiplicities and selection rules for the representations of $O(n, R)$ and $S(n+1)$ which subduce to the identity representation of C_{n+1} . This reduction we handle by character technique.

In the second step we characterize and construct an explicit set of orthonormal eigenmodes which span a unitary linear space of functions $\mathcal{L}^2$ for the harmonic analysis on the simplicial topological manifolds. The measure $d\mu$ on $\mathcal{L}^2$ in our analysis is taken over from the universal cover $S^{(n-1)}$. There are two alternative but equivalent approaches differing in the choice of the domain:

(I:) A given function $\mathcal{F}$ with domain $\mathcal{S}_0(n-1)$ can be extended to a C_{n+1} -periodic function on the universal cover $S^{(n-1)}$ and then analyzed exclusively in terms of a C_{n+1} -periodic basis.

(II:) By C_{n+1} -periodicity, this analysis can be collapsed entirely to the domain $\mathcal{S}_0(n-1)$, which forms a spherical tile of $S^{(n-1)}$ of volume $\text{vol}(S^{(n-1)})/(n+1)$, with the range of integration restricted to the topological manifold as domain. The expansion coefficients of a given function $\mathcal{F}$ with domain $\mathcal{S}_0(n)$ are found from the scalar products on $\mathcal{L}^2$ of $\mathcal{F}$ with the basis functions.

We focus on the choice (I) since the modification to (II) is straightforward. The main results of the following three sections are given in Tables after each section.

2 The 1-simplex $\mathcal{S}_0(1)$ on the circle S^1 .

As the simplest paradigm we treat here the spherical simplex $\mathcal{S}_0(1)$, with universal cover the unit circle S^1 according to Fig. 1, left.

2.1 The symmetric group $S(3)$.

Using orthogonal coordinates in the Euclidean plane E^2 , the symmetry group $S(3)$ of the regular triangle is generated by two Weyl reflections with matrices

$$(1, 2) : \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}, \quad (2, 3) : \begin{bmatrix} -\frac{1}{2} & \frac{1}{2}\sqrt{3} \\ \frac{1}{2}\sqrt{3} & \frac{1}{2} \end{bmatrix} \quad (6)$$

These matrices generate Young's orthogonal representation $D^{[21]}$ with basis vectors corresponding to the two Young tableaux

$$\begin{bmatrix} 1 & 3 \\ 2 & \end{bmatrix}, \quad \begin{bmatrix} 1 & 2 \\ 3 & \end{bmatrix}. \quad (7)$$

The two other irreducible representations of $S(3)$ are

$$\begin{aligned} D^{[3]} : D^{[3]}(1, 2) &= D^{[3]}(2, 3) = 1, \\ D^{[111]} : D^{[111]}(1, 2) &= D^{[111]}(2, 3) = -1, \end{aligned} \quad (8)$$

The character table of $S(3)$ in terms of classes, denoted by the cycle structures $k = (1)^3, (2)(1), (3)$ is given as **Table 2.1**.

2.2 The cyclic group C_3 .

The cyclic group as a subgroup $C_3 < S(3)$ is given in cycle notation by

$$C_3 = \{(1, 2, 3), (1, 2, 3)^2 = (3, 2, 1), (1, 2, 3)^3 = e\}. \quad (9)$$

In terms of the complex number $\lambda_3 = \exp(\frac{2\pi i}{3})$, the three irreducible representations $D^\alpha, \alpha = 0, 1, 2$ of C_3 coincide with their characters and are given for the group elements by **Table 2.2**.

2.3 The reduction $S(3) > C_3$.

The multiplicity $m(f, \alpha)$ of the subgroup irreducible representation D^α in the group irreducible representation D^f for $H < G$ is given from character technique by

$$m(f, \alpha) = \frac{1}{|H|} \sum_{k \in H} n(k) \chi^f(k) \overline{\chi}^\alpha(k). \quad (10)$$

Here $n(k)$ is the number of elements in class k of the subgroup H . We are interested only in the identity representation D^0 of $H = C_3$, collect the characters $\chi^f(k)$ of the group elements eq. 9 in the irreducible representations of $S(3)$ and $\chi^0(k) = D^0(k) = 1$, and compute the multiplicities $m(f, 0)$ from eq. 10 with $|C_3| = 3$, $n(k) = 1$ in **Table 2.3**.

As a result, the identity representation D^0 of C_3 is contained once in the irreducible representations $D^{[3]}, D^{[111]}$ of $S(3)$ but not in $D^{[21]}$.

2.4 The reduction $O(2, R) > S(3)$.

Now we turn to the group/subgroup pair $O(2) > S(3)$. The harmonic analysis on the circle S^1 is given by the complex Fourier series, spanned by the complete orthonormal set of functions of $z = \exp(i\phi)$, $0 \leq \phi < 2\pi$,

$$\begin{aligned} Y_m(z) &= \frac{1}{\sqrt{2\pi}} z^m, m = 0, \pm 1, \dots, \\ \int_0^{2\pi} d\phi \overline{Y_p}(z) Y_q(z) &= \delta_{pq}, \\ T_{(1,2)} Y_m(z) &= (-1)^m Y_{-m}. \end{aligned} \quad (11)$$

We include the reflection $(1, 2) : \phi \rightarrow \pi - \phi$ by introducing the new basis

$$Y_0, Y_{m,\epsilon} = \sqrt{\frac{1}{2}} [Y_m + \epsilon(-1)^m Y_{-m}], \epsilon = \pm 1, m = 1, 2, \dots \quad (12)$$

We are now ready to give the reduction of representations in $O(2) > S(3)$ from [5]. We introduce ν by

$$\nu : m \equiv \nu \bmod 3, \nu = 0, 1, 2. \quad (13)$$

Then the results from [5] pp. 255-7 can be combined with those from **Table 2.3** to yield the full reduction of representations for the groups eq. 5 in **Table 2.4**.

Table 2.4 determines the harmonic analysis on $\mathcal{S}_0(1)$ carried out on S^1 : Any function $\mathcal{F}$ with domain $\mathcal{S}_0(1)$ has a unique C_3 -periodic extension to S^1 and can be expanded into C_3 -periodic basis functions. The alternative would be to restrict the integration in eq. 11 to the sector $\mathcal{S}_0$ of angular range $2\pi/3$ and to reduce the volume from 2π to $2\pi/3$.

The C_3 -periodic basis is obtained from the irreducible representations of $O(2, R)$ by restriction to the values $\nu = 0, m \equiv 0 \bmod 3, \epsilon = \pm 1$. This restricts the irreducible representations of $S(3)$ to $f = [3], [111]$. Finally the representation of the point symmetry group $S(2)$ of the spherical simplex $\mathcal{S}_0(1)$ is given in terms of partitions f by

$$m = 0 : [2], m > 0, \epsilon = 1 : f = [2], \epsilon = -1 : f = [11]. \quad (14)$$

2.5 Tables for $O(2, R) > S(3) > C_3$.

Table 2.1. Characters for the 3 partitions f and classes k in cycle notation of the symmetric group $S(3)$.

$\chi^f(k)$	$(1)^3$	$(2)(1)$	(3)
$[3]$	1	1	1
$[21]$	2	0	-1
$[111]$	1	-1	1

(15)

Table 2.2. Characters for the 3 irreducible representations of C_3 with $\lambda_3 = \exp(\frac{2\pi i}{3})$.

χ^α	e	$(1, 2, 3)$	$(3, 2, 1)$
D^0	1	1	1
D^1	1	λ_3	λ_3^2
D^2	1	λ_3^2	λ_3

(16)

Table 2.3. Characters and multiplicity $m(f, 0)$ in the reduction of representations $S(3) > C_3$.

k	e	$(1, 2, 3)$	$(3, 2, 1)$	$m(f, 0)$
$\chi^{[3]}(k)$	1	1	1	1
$\chi^{[21]}(k)$	2	-1	-1	0
$\chi^{[111]}(k)$	1	1	1	1
$\chi^0(k)$	1	1	1	

(17)

Table 2.4. Reduction of irreducible representations for the groups $O(2, R) > S(3) > C_3$.

$O(2)$ (m, ϵ)	$S(3)$ f	C_3 $m(f, 0)$
$m = 0$	$[3]$	1
$m > 0, \nu = 0, \epsilon = 1$	$[3]$	1
$m > 0, \nu = 0, \epsilon = -1$	$[111]$	1
$m > 0, \nu = 1, \epsilon = \pm 1$	$[21]$	0
$m > 0, \nu = 2, \epsilon = \pm 1$	$[21]$	0

(18)

3 The 2-simplex $\mathcal{S}_0(2)$ on the sphere S^2 .

The 2-faces of the regular 3-simplex centrally projected to the sphere S^2 , Fig. 2, tiles it into 4 spherical triangles. The tiling has the symmetry $S(4)$ with the Coxeter diagram

$$S(4) : \circ - \circ - \circ. \quad (19)$$




Fig.2 The 3-simplex with vertices 1, 2, 3, 4 can be centrally projected to the sphere S^2 . The spherical triangle spanned by the vertices 1, 2, 3 on S^2 is taken as the simplicial topological manifold $\mathcal{S}_0(2)$.

The triangle obtained by dropping the vertex labelled 4 we take as the simplicial topological manifold $\mathcal{S}_0(2)$ seen on its universal cover S^2 . The generators $(1, 2), (2, 3), (3, 4)$ of $S(4)$ act as three reflection planes of the tetrahedron. The first two of them map $\mathcal{S}_0(2)$ into itself and generate the point group $S(3)$ of the simplicial manifold.

3.1 The group $S(4)$.

The characters of the irreducible representations of $S(4)$ are given from [3] p. 187 in **Table 3.1**.

3.2 The cyclic group C_4 .

The cyclic group C_4 is generated as

$$C_4 : \left\{ (1, 2, 3, 4), (1, 2, 3, 4)^2 = (1, 3)(2, 4), (1, 2, 3, 4)^3 = (4, 3, 2, 1), (1, 2, 3, 4)^4 = e \right\} \quad (20)$$

From these expressions it can be seen that the elements of C_4 generate all the cosets of the subgroup $S(3) < S(4)$. Hence their actions produce the full tiling of S^2 by triangles, are fixpoint-free and so fulfill all the properties required for $deck(\tilde{\mathcal{S}}_0(2))$. With $\lambda_4 = \exp(2\pi i/4) = i$, the 4 1-dimensional representations of C_4 are generated by

$$D^0(1, 2, 3, 4) = 1, D^1(1, 2, 3, 4) = i, D^2(1, 2, 3, 4) = i^2 = -1, D^3(1, 2, 3, 4) = i^3 = -i, \quad (21)$$

The characters of C_4 are given in **Table 3.2**.

3.3 The reduction $S(4) > C_4$.

We require the reduction of the representations f of $S(4)$ to the identity representation D^0 of C_4 . For this purpose we write the characters χ^f of these representations and $\chi^0 = D^0$ for the classes k of C_4 in the form of **Table 3.3** and compute the multiplicity $m(f, 0)$.

As a result, the identity representation D^0 of C_4 is contained once in the representations $f = [4], [22], [211]$ but not in $f = [31], [1111]$. Since the representations $f = [22], [211]$ are 2- and 3-dimensional, we must still find the 1-dimensional subspaces for the representation $D^0(C_4)$. This will be done by projection in subsection 3.5.1, 3.5.2.

3.4 The reduction $O(3, R) > S(4)$.

Next we turn to the harmonic analysis of $O(3, R)$ acting on the sphere S^2 . $O(3)$ is the direct product $SO(3, R) \times \mathcal{J}$, where $\mathcal{J}$ is the group generated by parity. Following [5], we characterize the irreducible representations by (l, κ) where $l = 0, 1, \dots$ is the integer angular momentum and $\kappa = \pm 1$ a parity label. The basis of the irreducible representation $D^l(SO(3, R))$ is spanned in spherical polar coordinates θ, ϕ by the spherical harmonics, [1] pp.19-25,

$$Y_{lm}(\theta, \phi), \quad -l \leq m \leq l, \quad (22)$$

$$\int \bar{Y}_{l'm'} Y_{lm} \sin(\theta) d\theta d\phi = \delta_{l'l} \delta_{m'm}.$$

which under the parity operation P transform as

$$PY_{lm} = (-1)^l Y_{lm}, \quad -l \leq m \leq l. \quad (23)$$

The parity therefore is

$$\kappa = (-1)^l. \quad (24)$$

In **Table 3.4** we reproduce from [5] pp. 259-60 the decomposition of irreducible representations for $(O(3, R) \times \mathcal{J}) > S(4)$ and combine it with the results from **Table 3.3**. Because of eq. 24, parity κ and l are correlated.

This Table can be extended by use of character technique given in [5].

Counting the total number of states up to $l = 4$ from **Table 3.4** one finds 25 basis states for the group $O(3, R)$. Only 7 of them contribute to the C_4 -periodic subset.

3.5 The explicit reduction $O(3, R) > S(4) > C_4$.

For higher representations (l, κ) the multiplicity $m((l, \kappa), f)$ of the partition f will take values larger than 1. In [5] we devised a way which allows for a complete orthogonal basis in this case. The idea is to introduce an additional hermitian operator, termed a generalized Casimir operator, as a polynomial in the components of the angular momentum operator, which for fixed (l, κ) by distinct eigenvalues separates repeated partitions f . We refer to [5] pp. 263-66 for the details.

We now determine the 1-dimensional subspaces for the partitions $f = [22], [211]$ belonging to the representation D^0 of C_4 .

3.5.1 The partition $f = [211]$.

Following [5], eq. (6.9), we choose particular coordinates with respect to the 4-simplex Fig. 2,

$$\begin{aligned} y_1 &= \frac{1}{2} [x_1 + x_2 - x_3 - x_4] \\ y_2 &= \frac{1}{2} [x_1 + x_3 - x_2 - x_4] \\ y_3 &= \frac{1}{2} [x_1 + x_4 - x_2 - x_3] \end{aligned} \quad (25)$$

These coordinates are convenient and stem from the analysis of the tetrahedral group T_d isomorphic to $S(4)$ [5] p. 258-68.

Expressed in these coordinates, the generators of $S(4)$ have the 3×3 matrix representation

$$D^{[31]'}(1, 2) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & -1 & 0 \end{bmatrix} \quad D^{[31]'}(2, 3) = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad D^{[31]'}(3, 4) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}. \quad (26)$$

which generate the representation $D^{[31]'}$ used in relation with the isomorphic tetrahedral group [5] p. 261. We use primes to distinguish this representation from the equivalent Young representation $D^{[31]}$ [3] pp. 225-6.

The representation with the associate partition $D^{[211]'}$ is obtained by multiplying each matrix for a generator in eq. 26 by (-1) , corresponding to the representation $D^{[111]}$. Within this representation we construct the matrix of the Coxeter element

$$D^{[211]'}(1, 2, 3, 4) = \begin{bmatrix} 0 & 0 & -1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}. \quad (27)$$

and by taking powers the representation for all 4 elements $h \in C_4$. The projection operator to the representation D^0 of C_4 becomes

$$P^{[211]',0} = \frac{1}{4} \sum_{h \in C_4} D^{[211]'}(h) = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}. \quad (28)$$

It picks the second basis function of $D^{[211]'}$ as the one which transforms according to D^0 .

Alternatively we describe the same representation $f = [211]$ in terms of the standard Young-Yamanouchi basis as given by Hamermesh [3] pp. 224-226. In contrast to the primed representation eq. 26 we label the basis vectors of this unprimed representation by three Yamanouchi symbols corresponding to three Young tableaux [3] pp 225-6:

$$1 : 3211, \begin{bmatrix} 1 & 2 \\ 3 & \\ 4 & \end{bmatrix}, 2 : 3121, \begin{bmatrix} 1 & 3 \\ 2 & \\ 4 & \end{bmatrix}, 3 : 1321, \begin{bmatrix} 1 & 4 \\ 2 & \\ 3 & \end{bmatrix}. \quad (29)$$

In terms of the canonical group/subgroup sequence $S(4) > S(3) > S(2)$ underlying the Young representation, the three Young basis vectors in eq. 29 correspond to the partitions $[21]$, $[21]$, $[111]$ of $S(3)$ in that order. We find the matrix representations of the reflection generators and of the generator of C_4 as

$$D^{[211]}(1, 2) : \begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{bmatrix} \quad D^{[211]}(2, 3) : \begin{bmatrix} -\frac{1}{2} & \frac{1}{2}\sqrt{3} & 0 \\ \frac{1}{2}\sqrt{3} & \frac{1}{2} & 0 \\ 0 & 0 & -1 \end{bmatrix} \quad (30)$$

$$D^{[211]}(3, 4) : \begin{bmatrix} -1 & 0 & 0 \\ 0 & -\frac{1}{3} & \sqrt{\frac{8}{9}} \\ 0 & \sqrt{\frac{8}{9}} & \frac{1}{3} \end{bmatrix}, D^{[211]}(1, 2, 3, 4) : \begin{bmatrix} \frac{1}{2} & -\frac{1}{6}\sqrt{3} & \sqrt{\frac{2}{3}} \\ \frac{1}{2}\sqrt{3} & \frac{1}{6} & -\frac{1}{3}\sqrt{2} \\ 0 & \frac{2}{3}\sqrt{2} & \frac{1}{3} \end{bmatrix}.$$

We can find the linear combination of the basis states belonging to $D^0(C_4)$ by constructing the eigenvector $\psi^{[211]0}$ of the representation matrix $D^{[211]}(1, 2, 3, 4)$ with eigenvalue 1. This eigenvector is found in terms of the Young tableau labels eq. 29 as

$$\psi^{[211],0} = (\sqrt{\frac{1}{2}}, \sqrt{\frac{1}{6}}, \sqrt{\frac{1}{3}}). \quad (31)$$

From the linear combination of the basis states with the coefficients eq. 31 we can draw a surprising conclusion. The Young representation basis eq. 29 displays explicitly the sub-partition of $S(3)$ as the part of the Young diagram occupied by the numbers 1, 2, 3. The basis vector eq. 31 of $D^0(C_4)$ in the Young representation $[211]$ of $S(4)$ is a generic mixture of the two inequivalent representations $[21], [111]$ of the group $S(3)$. Although we interpreted $S(3)$ as the point group of the simplicial manifold, the reduction to D^0 does not preserve a single representation of this subgroup. The cyclic group C_4 still provides four coset generators for the subgroup $S(3) < S(4)$, and so $S(4)$ is generated from all the products of elements from its subgroups $S(3), C_4$. Moreover our result eq. 31 shows that the use of the Young representation may not be optimal for the construction. Comparison of eq. 31 with the simpler result eq. 28 favours the primed tetrahedral form eq. 26 of the representation.

3.5.2 The partition $f = [22]$.

The second representation $D^{[22]}$ we treat in the Young orthogonal representation [3] p. 226 with the Young tableaux

$$1 : \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}, 2 : \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix}, \quad (32)$$

and with generators

$$D^{[22]}(1, 2) : \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, D^{[22]}(2, 3) : \begin{bmatrix} -\frac{1}{2} & -\frac{1}{2}\sqrt{3} \\ -\frac{1}{2}\sqrt{3} & \frac{1}{2} \end{bmatrix}, \quad (33)$$

$$D^{[22]}(3, 4) : \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, D^{[22]}(1, 2, 3, 4) : \begin{bmatrix} -\frac{1}{2} & \frac{1}{2}\sqrt{3} \\ \frac{1}{2}\sqrt{3} & \frac{1}{2} \end{bmatrix}.$$

By use of eq. 20 we find from the second line of eq. 33 the representation of the Coxeter element. The projection operator for the representation $D^0(C_4)$ becomes the matrix

$$P^{[22],0} = \frac{1}{4} \sum_{h \in C_4} D^{[22]}(h) = \frac{1}{4} \begin{bmatrix} 1 & \sqrt{3} \\ \sqrt{3} & 3 \end{bmatrix}. \quad (34)$$

and determines a unique linear combination in the Young tableaux,

$$\psi^{[22],0} = \left(\frac{1}{2}, \sqrt{\frac{3}{4}}\right) \quad (35)$$

as the basis for $D^0(C_4)$. The representation of the point group $S(3)$ of the simplicial manifold is fixed by the partition $f = [21]$.

This completes the group/subgroup basis construction of the harmonic analysis for a function $\mathcal{F}$ on the simplicial topological manifold $\mathcal{S}_0(2)$ when C_4 -periodically extended to S^2 . The new feature here is the appearance of multiplicity problems which require projections and the construction of appropriate operators. The C_4 -periodic basis for the harmonic analysis of $\mathcal{S}_0(2)$ on the sphere S^2 is characterized by the labels $((l, \kappa), f)$ of the irreducible representations of $O(3, R), S(4)$, with f restricted to allowed values.

3.6 Tables for $O(3, R) > S(4) > C_4$.

Table 3.1. Characters for the group $S(4)$.

$\chi^f(k)$	$(1)^4$	(4)	$(3)(1)$	$(2)^2$	$(2)(1)^2$
$[4]$	1	1	1	1	1
$[31]$	3	-1	0	-1	1
$[22]$	2	0	-1	2	0
$[211]$	3	1	0	-1	1
$[1111]$	1	-1	1	1	-1

(36)

Table 3.2. Characters for the cyclic group C_4 .

χ^α	e	$(1, 2, 3, 4)$	$(1, 3)(2, 4)$	$(4, 3, 2, 1)$
D^0	1	1	1	1
D^1	1	i	-1	$-i$
D^2	1	-1	1	-1
D^3	1	$-i$	-1	i

(37)

Table 3.3. Characters and multiplicities in the reduction $S(4) > C_4$.

k	e	$(1, 2, 3, 4)$	$(1, 3)(2, 4)$	$(4, 3, 2, 1)$	$m(f, 0)$
$\chi^{[4]}(k)$	1	1	1	1	1
$\chi^{[31]}(k)$	3	-1	-1	-1	0
$\chi^{[22]}(k)$	2	0	0	2	1
$\chi^{[211]}(k)$	3	1	1	-1	1
$\chi^{[1111]}(k)$	1	-1	-1	1	0
$\chi^0(k)$	1	1	1	1	

(38)

Table 3.4. The multiplicities $m((l, \kappa), f)$ of representations D^f of $S(4)$ and $m((\lambda, \kappa), 0)$ of C_4 -periodic eigenmodes in the general reduction $O(3) > S(4) > C_4$ for $(l, \kappa), l \leq 4$. Because of eq. 24, only the pairs (l, κ) with $\kappa = (-1)^l$ contribute to the C_4 -periodic modes.

$(l, \kappa), f :$	[4]	[31]	[22]	[211]	[1111]	$m((l, \kappa), 0)$
$(0, 1)$	1	0	0	0	0	1
$(1, -1)$	0	1	0	0	0	0
$(2, 1)$	0	1	1	0	0	1
$(3, -1)$	1	1	0	1	0	2
$(4, 1)$	1	1	1	1	0	3

(39)

4 The tetrahedral 3-simplex $\mathcal{S}_0(3)$ on the sphere S^3 .

We consider the 3-sphere $S^3 < E^4$ and an inscribed regular 4-simplex with its vertices enumerated as 1, 2, 3, 4, 5. The full point symmetry of the 4-simplex is $S(5)$. Central projection of the 3-faces of this simplex to S^3 yields a tiling with 5 tetrahedral tiles. We choose the tetrahedron obtained by dropping the vertex 5 as the simplicial manifold $\mathcal{S}_0(3)$. Its internal point symmetry group is $S(4)$. The homotopy group $\pi_1(\mathcal{S}_0(3))$ of the Platonic tetrahedron is described by Everitt in [2] by a graph algorithm. Its prime dimension 5 identifies it and the group of deck transformations as the cyclic group C_5 . In the next subsections we implement the group/subgroup analysis in analogy to the previous sections with the goal to characterize the harmonic analysis on $\mathcal{S}_0(3)$.

4.1 The reduction $S(5) > C_5$.

The characters of the irreducible representations of $S(5)$ for all 7 classes and irreducible representations from [3] p. 276 are given in **Table 4.1**.

We shall need the characters of Table 4.1 for the reduction of the rotation group.

The cyclic group C_5 has the elements

$$g = (1, 2, 3, 4, 5), g^2 = (1, 3, 5, 2, 4), g^3 = (1, 4, 2, 5, 3), g^4 = (1, 5, 4, 3, 2), g^5 = e. \quad (40)$$

They belong to the classes (4)(1) or (5) of $S(5)$. The computation of the multiplicity $m(f, 0)$ of the identity representation $D^0(C_5)$ is straightforward and we include it in the last column of **Table 4.1**. The representation D^0 is contained once in the representations $f = [5], [11111], [32], [221]$, twice in the representation $f = [311]$, but not in the representations $f = [41], f = [2111]$. Again we must determine the corresponding subspaces for the first set of representations.

4.2 $O(4, R)$ and Weyl reflections acting on S^3 .

We shall adopt from [6] the coordinate description of the sphere S^3 , equivalent to an element $u \in SU(2, C)$, and relate it to the vector notation $x = (x_0, x_1, x_2, x_3)$ in E^4 by

$$u = \begin{bmatrix} z_1 & z_2 \\ -\bar{z}_2 & \bar{z}_1 \end{bmatrix}, \det(u) = z_1 \bar{z}_1 + z_2 \bar{z}_2 = 1, \quad (41)$$

$$z_1 = x_0 - ix_3, \quad z_2 = -(x_2 + ix_1).$$

We use the isomorphism $SO(4, R) \sim (SU^l(2, C) \times SU^r(2, C))/Z_2$ with the action defined as

$$(SU^l(2, C) \times SU^r(2, C)) \times S^3 \rightarrow S^3, \quad (42)$$

$$(g_l, g_r) \times u \rightarrow (g_l)^{-1} u g_r.$$

Here Z_2 denotes the subgroup generated by the action of $(-e, -e) \in (SU^l(2, C) \times SU^r(2, C))$ in eq. 42 which preserves any point $u \in S^3$. We stress that the half-angular parameters here must be used in their full range appropriate for $SU(2)$. The spherical harmonics on S^3 are shown in [6] to be equal to the Wigner representation functions D^j of $SU(2)$ eq. 85,

$$Y_{m_1 m_2}^j(u) = D_{m_1 m_2}^j(u), \quad j = 0, 1/2, 1, 3/2, \dots, -j \leq m_i \leq j, \quad i = 1, 2, \quad (43)$$

$$\int \bar{D}_{m'_1 m'_2}^{j'}(u) D_{m_1 m_2}^j(u) d\mu(u) = \delta_{j', j} \delta_{m'_1, m_1} \delta_{m'_2, m_2}.$$

The measure $d\mu(u)$ can be expressed by three Euler angle parameters [1] pp. 62-3. We denote the irreducible representation obtained by the action of rotations and Weyl reflections on the spherical harmonics eq. 43 as $D^{(j, j)}$. In the appendix we summarize some properties of the Wigner D^j functions. In terms of the complex numbers $(z_1, z_2, \bar{z}_1, \bar{z}_2)$ in eq. 41, the spherical harmonics are homogeneous polynomials of degree $(2j)$, see eq. 85. Under the action of rotations they transform, due to their

representations properties, irreducibly with a representation $D^{(j,j)}$ given by direct product matrices,

$$\begin{aligned}
(T_{(g_l, g_r)} D_{m_1 m_2}^j)(u) &= D_{m_1, m_2}^j(g_l^{-1} u g_r) \\
&= \sum_{m'_1, m'_2} D_{m'_1 m'_2}^j(u) \left[D_{m_1 m'_1}^j(g_l^{-1}) D_{m'_2 m_2}^j(g_r) \right], \\
D_{m'_1 m'_2, m_1 m_2}^{(j,j)}(g_l, g_r) &= \langle (j, j)(m'_1 m'_2) | T_{(g_l, g_r)} | (j, j)(m_1 m_2) \rangle \\
&= D_{m_1 m'_1}^j(g_l^{-1}) D_{m'_2 m_2}^j(g_r)
\end{aligned} \tag{44}$$

where finally we passed for convenience to a bracket notation. The defining action of $SO(4, R)$ on E^4 by real 4×4 rotation matrices is included in this notation as the representation $D^{(\frac{1}{2}, \frac{1}{2})}$.

For the character of the irreducible representation $D^{(j,j)}$ of a rotation it follows from eq. 44 that it is given by

$$\chi^{(j,j)}(T_{(g_l, g_r)}) = \text{Trace}(D^j(g_l^{-1}) \times D^j(g_r)) = \chi^j(g_l^{-1}) \chi^j(g_r), \tag{45}$$

in terms of pairs of characters $\chi^j(g)$ of $SU(2, C)$. The dimension of $D^{(j,j)}$ from eq. 45 is $(\chi^j(e))^2 = (2j+1)^2$.

To determine the action of the symmetric group $S(5)$ on S^3 we need the action of a Weyl reflection eq. 4 in the coordinate description eq. 41 and at the same time must extend the action from $SO(4, R)$ to $O(4, R)$. We start with the Weyl reflection for the particular Weyl vector $a_0 = (1, 0, 0, 0)$ which acts as

$$(W_{a_0} \times (x_0, x_1, x_2, x_3)) \rightarrow (-x_0, x_1, x_2, x_3). \tag{46}$$

Inserting this reflection into the complex coordinates of u from equation 41 we find that the image of u under W_{a_0} can be rewritten in matrix notation as

$$W_{a_0}(u) = -u^\dagger. \tag{47}$$

In addition we need the representations and characters for reflections from $O(4, R)$. For $u \in SU(2, C)$ we need the properties eqs. 83 and 87. This equation and the homogeneity eq. 88 of degree $2j$ allows us to rewrite the Weyl operator T_{a_0} acting on a spherical harmonic eq.43 as

$$\begin{aligned}
(T_{a_0} D_{m_1 m_2}^j)(u) &= D_{m_1 m_2}^j(-u^\dagger) \\
&= (-1)^{2j} D_{m_2 m_1}^j(q^{-1} u q) = (-1)^{2j} (T_{(q, q)} D_{m_2 m_1}^j)(u).
\end{aligned} \tag{48}$$

We used the homogeneity eq. 88 to extract the phase factor $(-1)^{2j}$. With respect to an action eq. 44 of $SO(4, R)$ following the reflection $W_{a_0}(u)$, we get from eq. 48

$$((T_{(g_l, g_r)} T_{a_0}) D_{m_1, m_2}^j)(u) = (-1)^{2j} ((T_{(g_l, g_r)} T_{(q, q)}) D_{m_2, m_1}^j)(u) \tag{49}$$

For the Weyl operator T_{a_0} eq. 48 we find the involutive and conjugation properties

$$\begin{aligned} T_{a_0} T_{a_0} &= I, \\ T_{a_0} T_{(g_l, g_r)} T_{a_0} &= T_{(g_r, g_l)} \end{aligned} \quad (50)$$

The Weyl vector a_0 in the coordinates eq. 41 corresponds to $u = e$. For a general unit Weyl vector a we can choose a rotation $R \in SO(4, R)$ such that $a = Ra_0$. Then by the general operator relation between Weyl reflections and rotations

$$W_a = W_{Ra_0} = RW_{a_0}R^{-1}, \quad (51)$$

the general Weyl reflection W_a can be expressed in terms of the particular reflection W_{a_0} . It remains to construct a rotation R . In terms of the coordinates eq. 41 we note that

$$((e, v) \times e) = v \quad (52)$$

which means that the element $(e, v) \in (SU^l(2, C) \times SU^r(2, C))$ applied to a_0 gives the most general point $a \in S^3$ and so provides the rotation R .

With this choice of R , the action of a general Weyl reflection operator on a spherical harmonic can be written by use of eq. 50 as

$$T_a = T_{(e, v_a^{-1})} T_{a_0} T_{(e, v_a)} = T_{(v_a, v_a^{-1})} T_{a_0} \quad (53)$$

Here the matrix v_a is obtained from the Weyl vector a by inserting the vector components of $x = a$ into the matrix eq. 41. From the factorization eq. 53 of the general Weyl reflection operator and from eq. 44 we find the matrix representation of T_a as

$$\begin{aligned} &\langle (j, j)(m'_1 m'_2) | T_{(v_a, v_a^{-1})} T_{a_0} | (j, j)(m_1 m_2) \rangle \\ &= (-1)^{2j} \langle (j, j)(m'_1 m'_2) | T_{(v_a, v_a^{-1})} T_{(q, q)} | (j, j)(m_2 m_1) \rangle \\ &= (-1)^{2j} D_{m_2 m'_1}^j(q^{-1} v_a^{-1}) D_{m'_2 m_1}^j(v_a^{-1} q). \end{aligned} \quad (54)$$

Here we used the complex conjugation property $\bar{u} = q^{-1} u q$ of $SU(2, C)$, eq. 83. The trace of this representation becomes

$$\chi^{(j, j)}(T_a) = (-1)^{2j} \chi^j(q^{-1} v_a^{-1} (-q) (v_a^{-1})^T) = \chi^j(\overline{v_a^{-1}} (v_a^{-1})^T) = \chi^j(e) = 2j + 1. \quad (55)$$

This result is independent of v as expected from eq. 53 since all Weyl operators by eq. 51 are conjugate to T_{a_0} and the trace is independent of conjugations.

All operators which are not pure rotations can be written with the help of eq. 50 in the form

$$T = T_{(g_l, g_r)} T_{a_0}. \quad (56)$$

The matrix elements of this operator are given similar to eq. 54 by

$$\begin{aligned} &(-1)^{2j} \langle (j, j)(m'_1 m'_2) | T_{(g_l, g_r)} | (j, j)(m_2 m_1) \rangle \\ &= (-1)^{2j} D_{m_2 m'_1}^j(q^{-1} g_l^{-1}) D_{m'_2 m_1}^j(g_r q). \end{aligned} \quad (57)$$

with q given in eq. 83. It follows from eq. 57 that the character of this operator in the representation $D^{(j,j)}$ is given by

$$\begin{aligned}\chi^{(j,j)}(T_{(g_l, g_r)} T_{a_0}) &= (-1)^{2j} \sum_{m_1 m_2} D_{m_2 m_1}^j(q^{-1} g_l^{-1}) D_{m_2 m_1}^j(g_r q) \\ &= \chi^j(\overline{g_l^\dagger} g_r^T) = \chi^j((g_r g_l)^T) = \chi^j(g_r g_l)\end{aligned}\quad (58)$$

Here we used from eq. 83 $q^T = (-q)$, which leads to the cancellation of the phase factor $(-1)^{2j}$, and $q^{-1} g_l^{-1} q = q^{-1} g_l^\dagger q = g_l^T$.

4.3 The representations of $S(5)$.

From the Coxeter diagram of $S(5)$ we can construct in E^4 a set of 4 Weyl vectors associated with the 4 generators. The scalar products must form, [4] pp. 108-10, the matrix

$$\langle a_i, a_j \rangle = \begin{bmatrix} 1 & \frac{1}{2} & 0 & 0 \\ \frac{1}{2} & 1 & \frac{1}{2} & 0 \\ 0 & \frac{1}{2} & 1 & \frac{1}{2} \\ 0 & 0 & \frac{1}{2} & 1 \end{bmatrix} \quad (59)$$

Here the off-diagonal scalar products are determined from the Coxeter diagram by $\langle a_i, a_{i+1} \rangle = \cos(\pi/3) = \frac{1}{2}$, $i = 1, 2, 3$. We fulfill eq. 59 by the choice of the four Weyl vectors given in **Table 4.2**.

From eq. 53 we compute for each Weyl vector $w_i, i = 1, \dots, 4$ the matrices v_i which appear in the reflection operator and include them in the last column of **Table 4.2**. From the generators of $S(5)$ we can construct representatives g_k of the seven classes k appearing in **Table 4.2** and express them as products of the Weyl reflection operators $W_i := W_{a_i}$ in **Table 4.3**.

For products of up to 4 Weyl reflection operators eq. 53 we find with eq. 50

$$\begin{aligned}T_a &= T_{(v_a, v_a^{-1})} T_{a_0}, \\ T_b T_a &= T_{(v_b v_a^{-1}, v_b^{-1} v_a)}, \\ T_c T_b T_a &= T_{(v_c v_b^{-1} v_a, v_c^{-1} v_b v_a^{-1})} T_{a_0}, \\ T_d T_c T_b T_a &= T_{(v_d v_c^{-1} v_b v_a^{-1}, v_d^{-1} v_c v_b^{-1} v_a)}.\end{aligned}\quad (60)$$

Here products with an even number of Weyl operators are rotations, those with an odd number become a product of a rotation with T_{a_0} .

4.4 Multiplicity in the reduction $O(4, R) > S(5) > C_5$.

To write explicitly the action of the class representatives from **Table 4.3** we must convert the corresponding product of Weyl operators for each class representative

with the help of eq. 60 into an explicit action of $SU^l(2, C) \times SU^r(2, C)$ on u or on $(-u^\dagger)$ respectively. Once we have computed these actions, the characters in the representation $D^{(j,j)}$ are found by use of eqs. 44, 58 respectively. It turns out that for five classes $k \in S(5)$ the characters $\chi^{(j,j)}(k)$ are periodic with periods 2, 3, 4, 5 respectively wrt. $(2j)$.

The results and corresponding recursion relations are given in **Tables 4.7, 4.8**.

The real scalar product

$$m((j, j), f) = \frac{1}{5!} \sum_{k \in S(5)} n(k) \chi^{(j,j)}(k) \chi^f(k), \quad (61)$$

with $n(k)$ the number of elements in class $k \in S(5)$, then yields the multiplicity $m((j, j), f)$ in the reduction $O(4, R) > S(5)$. In **Table 4.9** we write down the multiplicities computed according to eq. 61 for the reduction of representations in $O(4, R) > S(5)$. The multiplicity of C_5 -periodic modes of $O(4, R)$ is then given by

$$m((j, j), 0) = \sum_f m((j, j), f) m(f, 0). \quad (62)$$

in the last column of **Table 4.9**.

Of the 506 harmonic polynomials for $O(4, R)$ up to degree $2j = 10$, 101 are C_5 -periodic eigenmodes of $\mathcal{S}_0(3)$. and so demonstrate on average the topological selection rules for the simplicial manifold $\mathcal{S}^0(3)$. There are other specific selection rules like the absence of C_5 -periodic eigenmodes for $2j = 1$. By use of eq. 61 and the characters given in **Tables 4.7, 4.8**, **Table 4.9** can be extended to any value $(2j) > 10$. We note another recursion relation for the multiplicity $m((j, j), f)$: for all except the first two classes, the characters $\chi^{(j,j)}(k)$ from **Tables 4.7, 4.8** obey

$$k = (3)(1)^2, (2)^2(1), (3)(2), (4)(1), (5) : \chi^{(j+30, j+30)}(k) = \chi^{(j,j)}(k). \quad (63)$$

Using in eq. 61 the characters for the first two classes from **Table 4.9** and for the others the periodicity eq. 63 one finds for the multiplicity the recursion relation

$$m((j + 30, j + 30), f) = m((j, j), f) + (2j + 36). \quad (64)$$

4.5 The explicit reduction $S(5) > C_5$.

For the partitions $f = [32], [221], [311]$ we now compute the C_5 -periodic states ψ^{f0} as linear combinations of the basis states ϕ_j in the Young orthogonal representation. For each representation D^f which admits D^0 , they are given as the single or at most two eigenvectors of eigenvalue 1 for the Coxeter element generating C_5 in the

representations spaces for those partitions which from **Table 4.1** reduce to D^0 . We give as a vector the coefficients ψ_j^{f0} in the linear combinations

$$\psi^{f0} = \sum_j \phi_j \psi_j^{f0}. \quad (65)$$

The representation matrices for the generators of $S(5)$ and C_5 and the coefficients in eq. 65 are given in **Tables 4.4-4.6**.

4.6 Tables for $O(4, R) > S(5) > C_5$.

Table 4.1. Characters for the group $S(5)$ for irreducible representations D^f with partition f , classes k in cycle notation, number $n(k)$ of elements in class k , and multiplicity $m(f, 0)$ of the identity representation $D^0(C_5)$.

$\chi^f(k)$	$(1)^5$	$(2)(1)^3$	$(2)^2(1)$	$(3)(1)^2$	$(3)(2)$	$(4)(1)$	(5)	$m(f, 0)$
$n(k)$	1	10	15	20	20	30	24	
[5]	1	1	1	1	1	1	1	1
[11111]	1	-1	1	1	-1	-1	1	1
[41]	4	2	0	1	-1	0	-1	0
[2111]	4	-2	0	1	1	0	-1	0
[32]	5	1	1	-1	1	-1	0	1
[221]	5	-1	1	-1	-1	1	0	1
[311]	6	0	-2	0	0	0	1	2

Table 4.2. Weyl vectors a_i and matrices v_i in the Weyl operators eq. 53 for the generators $(i, i+1)$ of the group $S(5)$.

i	$(i, i+1)$	a_i	v_i
1	(1, 2)	$(0, 0, 0, 1)$	$\begin{bmatrix} -i & 0 \\ 0 & i \end{bmatrix}$
2	(2, 3)	$(0, 0, \sqrt{\frac{3}{4}}, \frac{1}{2})$	$\begin{bmatrix} -\frac{i}{2} & -\frac{1}{2}\sqrt{3} \\ \frac{1}{2}\sqrt{3} & \frac{i}{2} \end{bmatrix}$
3	(3, 4)	$(0, \sqrt{\frac{2}{3}}, \sqrt{\frac{1}{3}}, 0)$	$\begin{bmatrix} 0 & -(\sqrt{\frac{1}{3}} + i\sqrt{\frac{2}{3}}) \\ (\sqrt{\frac{1}{3}} - i\sqrt{\frac{2}{3}}) & 0 \end{bmatrix}$
4	(4, 5)	$(\sqrt{\frac{5}{8}}, \sqrt{\frac{3}{8}}, 0, 0)$	$\begin{bmatrix} \sqrt{\frac{5}{8}} & -i\sqrt{\frac{3}{8}} \\ -i\sqrt{\frac{3}{8}} & \sqrt{\frac{5}{8}} \end{bmatrix}$

Table 4.3. Representatives for the seven classes k of $S(5)$ in terms of Weyl reflections $W_i := W_{a_i}$, and matrices $g_l, g_r, g_r g_l$ appearing in eqs. 45, 56, 58.

$k :$	$(1)^5$	$(2)(1)^3$	$(2)^2(1)$	$(3)(1)^2$
g_k	e	$(1, 2)$	$(1, 2)(3, 4)$	$(1, 2)(2, 3)$
	I	W_1	$W_1 W_3$	$W_1 W_2$
g_l			$\begin{bmatrix} 0 & \frac{\sqrt{2}-i}{\sqrt{3}} \\ -\frac{\sqrt{2}+i}{\sqrt{3}} & 0 \end{bmatrix}$	$\begin{bmatrix} \frac{1}{2} & -\frac{i\sqrt{3}}{2} \\ -\frac{i\sqrt{3}}{2} & \frac{1}{2} \end{bmatrix}$
g_r			$\begin{bmatrix} 0 & \frac{\sqrt{2}-i}{\sqrt{3}} \\ -\frac{\sqrt{2}+i}{\sqrt{3}} & 0 \end{bmatrix}$	$\begin{bmatrix} \frac{1}{2} & -\frac{i\sqrt{3}}{2} \\ -\frac{i\sqrt{3}}{2} & \frac{1}{2} \end{bmatrix}$
$g_r g_l$				

(68)

$k :$	$(3)(2)$	$(4)(1)$	(5)
g_k	$(1, 2)(2, 3)(4, 5)$	$(1, 2)(2, 3)(3, 4)$	$(1, 2)(2, 3)(3, 4)(4, 5)$
	$W_1 W_2 W_4$	$W_1 W_2 W_3$	$W_1 W_2 W_3 W_4$
g_l			$\begin{bmatrix} \frac{2-2\sqrt{5}-i(\sqrt{2}+\sqrt{10})}{8} & \frac{3\sqrt{2}-\sqrt{10}+i(-6-2\sqrt{5})}{8\sqrt{3}} \\ \frac{-3\sqrt{2}+\sqrt{10}+i(-6-2\sqrt{5})}{8\sqrt{3}} & \frac{2-2\sqrt{5}+i(\sqrt{2}+\sqrt{10})}{8} \end{bmatrix}$
g_r			$\begin{bmatrix} \frac{2+2\sqrt{5}+i(-\sqrt{2}+\sqrt{10})}{8} & \frac{3\sqrt{2}+\sqrt{10}+i(-6+2\sqrt{5})}{8\sqrt{3}} \\ \frac{-3\sqrt{2}-\sqrt{10}+i(-6+2\sqrt{5})}{8\sqrt{3}} & \frac{2+2\sqrt{5}+i(\sqrt{2}-\sqrt{10})}{8} \end{bmatrix}$
$g_r g_l$	$\begin{bmatrix} -\frac{1}{2} & -\frac{i\sqrt{3}}{2} \\ -\frac{i\sqrt{3}}{2} & -\frac{1}{2} \end{bmatrix}$	$\begin{bmatrix} -\frac{i}{\sqrt{2}} & -\frac{\sqrt{2}+2i}{2\sqrt{3}} \\ \frac{\sqrt{2}-2i}{2\sqrt{3}} & \frac{i}{\sqrt{2}} \end{bmatrix}$	

(69)

Table 4.4. Periodic state for the partition $f = [32]$. Young tableau basis:

$$\begin{aligned}
 1 &= \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & \end{bmatrix}, 2 = \begin{bmatrix} 1 & 3 & 4 \\ 2 & 5 & \end{bmatrix}, 3 = \begin{bmatrix} 1 & 2 & 4 \\ 3 & 5 & \end{bmatrix}, \\
 4 &= \begin{bmatrix} 1 & 3 & 5 \\ 2 & 4 & \end{bmatrix}, 5 = \begin{bmatrix} 1 & 2 & 5 \\ 3 & 4 & \end{bmatrix}.
 \end{aligned}
 \tag{70}$$

Representation $D^{[32]}$ for generators from [3] pp. 227-8:

$$D^{[32]}(1, 2) = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}, \quad D^{[32]}(2, 3) = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & \frac{1}{2} & \sqrt{\frac{3}{4}} & 0 & 0 \\ 0 & \sqrt{\frac{3}{4}} & -\frac{1}{2} & 0 & 0 \\ 0 & 0 & 0 & \frac{1}{2} & \sqrt{\frac{3}{4}} \\ 0 & 0 & 0 & \sqrt{\frac{3}{4}} & -\frac{1}{2} \end{bmatrix} \tag{71}$$

$$D^{[32]}(3, 4) = \begin{bmatrix} -\frac{1}{3} & 0 & \sqrt{\frac{8}{9}} & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ \sqrt{\frac{8}{9}} & 0 & \frac{1}{3} & 0 & 0 \\ 0 & 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}, \quad D^{[32]}(4, 5) = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & -\frac{1}{2} & 0 & \sqrt{\frac{3}{4}} & 0 \\ 0 & 0 & -\frac{1}{2} & 0 & \sqrt{\frac{3}{4}} \\ 0 & \sqrt{\frac{3}{4}} & 0 & \frac{1}{2} & 0 \\ 0 & 0 & \sqrt{\frac{3}{4}} & 0 & \frac{1}{2} \end{bmatrix}.$$

The matrix $D^{[32]}(4, 5)$ given in [3] p. 228 has an error and was replaced.
Representation of Coxeter element of C_5 :

$$D^{[32]}((1, 2)(2, 3)(3, 4)(4, 5)) = \begin{bmatrix} -\frac{1}{3} & 0 & -\sqrt{\frac{2}{9}} & 0 & \sqrt{\frac{2}{3}} \\ -\sqrt{\frac{2}{3}} & \frac{1}{4} & \sqrt{\frac{1}{48}} & -\sqrt{\frac{3}{16}} & -\frac{1}{4} \\ -\sqrt{\frac{2}{9}} & -\sqrt{\frac{3}{16}} & \frac{1}{12} & \frac{3}{4} & -\sqrt{\frac{1}{48}} \\ 0 & \sqrt{\frac{3}{16}} & -\frac{3}{4} & \frac{1}{4} & -\sqrt{\frac{3}{16}} \\ 0 & -\frac{3}{4} & -\sqrt{\frac{3}{16}} & -\sqrt{\frac{3}{16}} & -\frac{1}{4} \end{bmatrix} \quad (72)$$

Eigenvector of Coxeter element for the single eigenvalue 1 eq. 65:

$$\psi^{[32]0} = (\sqrt{\frac{2}{3}}, -1, -\sqrt{\frac{1}{3}}, -\sqrt{\frac{1}{3}}, 1). \quad (73)$$

Table 4.5. Periodic state for the partition $f = [221]$. The Young tableaux of the basis are the mirror images of those for the associate partition $[32]$ eq. 70. The representation matrices can then be computed from those of $D^{[32]}$ eq. 71.

Representation of the Coxeter element:

$$D^{[221]}((1, 2)(2, 3)(3, 4)(4, 5)) = \begin{bmatrix} -\frac{1}{3} & 0 & \sqrt{\frac{2}{9}} & 0 & \sqrt{\frac{2}{3}} \\ -\sqrt{\frac{2}{3}} & \frac{1}{4} & -\sqrt{\frac{1}{48}} & \sqrt{\frac{3}{16}} & -\frac{1}{4} \\ \sqrt{\frac{2}{9}} & \sqrt{\frac{3}{16}} & \frac{1}{12} & \frac{3}{4} & \sqrt{\frac{1}{48}} \\ 0 & -\sqrt{\frac{3}{16}} & -\frac{3}{4} & \frac{1}{4} & \sqrt{\frac{3}{16}} \\ 0 & -\frac{3}{4} & \sqrt{\frac{3}{16}} & \sqrt{\frac{3}{16}} & -\frac{1}{4} \end{bmatrix} \quad (74)$$

Eigenvector of the Coxeter element for the eigenvalue 1:

$$\psi^{[221]0} = (\sqrt{\frac{2}{3}}, -1, \sqrt{\frac{1}{3}}, \sqrt{\frac{1}{3}}, 1). \quad (75)$$

Table 4.6. Periodic state for the partition $f = [311]$ in eq. 65:

Young tableau basis:

$$1 = \begin{bmatrix} 1 & 2 & 3 \\ 4 & & \\ 5 & & \end{bmatrix}, \quad 2 = \begin{bmatrix} 1 & 2 & 4 \\ 3 & & \\ 5 & & \end{bmatrix}, \quad 3 = \begin{bmatrix} 1 & 3 & 4 \\ 2 & & \\ 5 & & \end{bmatrix}, \quad (76)$$

$$4 = \begin{bmatrix} 1 & 2 & 5 \\ 3 & & \\ 4 & & \end{bmatrix}, \quad 5 = \begin{bmatrix} 1 & 3 & 5 \\ 2 & & \\ 4 & & \end{bmatrix}, \quad 6 = \begin{bmatrix} 1 & 4 & 5 \\ 2 & & \\ 3 & & \end{bmatrix},$$

The representation matrices of the generators are given in [3] pp. 229-30. Computed matrix representation of the Coxeter element:

$$D^{[311]}((1, 2)(2, 3)(3, 4)(4, 5)) = \begin{bmatrix} \frac{1}{3} & -\sqrt{\frac{1}{18}} & 0 & \sqrt{\frac{5}{6}} & 0 & 0 \\ \sqrt{\frac{2}{9}} & \frac{1}{24} & -\sqrt{\frac{3}{64}} & -\sqrt{\frac{5}{192}} & \sqrt{\frac{45}{64}} & 0 \\ \sqrt{\frac{2}{3}} & \sqrt{\frac{1}{192}} & \frac{1}{8} & -\sqrt{\frac{5}{64}} & -\sqrt{\frac{15}{64}} & 0 \\ 0 & \sqrt{\frac{15}{64}} & -\sqrt{\frac{5}{64}} & \frac{1}{8} & -\sqrt{\frac{1}{192}} & \sqrt{\frac{2}{3}} \\ 0 & \sqrt{\frac{45}{64}} & \sqrt{\frac{5}{192}} & \sqrt{\frac{3}{64}} & \frac{1}{24} & -\sqrt{\frac{2}{9}} \\ 0 & 0 & \sqrt{\frac{5}{6}} & 0 & \sqrt{\frac{1}{18}} & \frac{1}{3} \end{bmatrix} \quad (77)$$

Two eigenvectors in eq. 65 for eigenvalue 1 of eq. 77:

$$\begin{aligned} q^{[311]0,1} &= (\sqrt{\frac{49}{45}}, \sqrt{\frac{2}{45}}, \sqrt{\frac{8}{15}}, \sqrt{\frac{2}{3}}, 0, 1), \\ q^{[311]0,2} &= (\sqrt{\frac{8}{45}}, \sqrt{\frac{49}{45}}, -\sqrt{\frac{1}{15}}, \sqrt{\frac{1}{3}}, 1, 0), \end{aligned} \quad (78)$$

Table 4.7. The angles ϕ and characters $\chi^{(j,j)}(k)$ for the classes $k \in S(5)$ and the representations $D^{(j,j)}$ of $O(4, R)$ with $(2j) = 0, 1, 2, 3, 4, 5$. General expressions and recursion relations are given in **Table 4.8**.

k	ϕ	$\chi^{(j,j)}(k), (2j) :$	0	1	2	3	4	5
$(1)^5$	$\phi(g_l)/2 = \phi(g_r)/2 = 0$		1	4	9	16	25	36
$(2)(1)^3$	$\phi(g_r g_l)/2 = 0$		1	2	3	4	5	6
$(3)(1)^2$	$\phi(g_l)/2 = \phi(g_r)/2 = \pi/3$		1	1	0	1	1	0
$(2)^2(1)$	$\phi(g_l)/2 = \phi(g_r)/2 = \pi/2$		1	0	1	0	1	0
$(3)(2)$	$\phi(g_r g_l)/2 = 2\pi/3$		1	-1	0	1	-1	0
$(4)(1)$	$\phi(g_r g_l)/2 = \pi/2$		1	0	-1	0	1	0
(5)	$\phi(g_l)/2 = 2\pi/10, \phi(g_r)/2 = 6\pi/10$		1	-1	-1	1	0	1

(79)

Table 4.8 General expressions and recursion relations for the characters $\chi^{(j,j)}(k)$ as functions of classes k of $S(5)$. The initial values are given in **Table 4.7**.

k	$\chi^{(j,j)}(k)$
$(1)^5$	$\chi^{(j,j)} = (2j+1)^2$
$(2)(1)^3$	$\chi^{(j,j)} = (2j+1)$
$(3)(1)^2$	$\chi^{(j+3/2, j+3/2)} = \chi^{(j,j)}$
$(2)^2(1)$	$\chi^{(j+1, j+1)} = \chi^{(j,j)}$
$(3)(2)$	$\chi^{(j+3/2, j+3/2)} = \chi^{(j,j)}$
$(4)(1)$	$\chi^{(j+2, j+2)} = \chi^{(j,j)}$
(5)	$\chi^{(j+5/2, j+5/2)} = \chi^{(j,j)}$

(80)

Table 4.9. Multiplicities $m((j, j), f)$ in the reduction of representations $D^{(j,j)} = \sum_j m((j, j), f) D^f$ from $O(4, R)$ to $S(5)$ as function of $(2j) = 0, \dots, 10$ and of all partitions f . The numbers $m((j, j), 0)$ in the last column denotes the total number of C_5 -periodic modes for fixed $(2j)$, $\nu_0(f)$ in the last row the total number for a fixed partition f up to $(2j) = 10$.

f	[5]	[11111]	[41]	[2111]	[32]	[221]	[311]	$m((j, j), 0)$
$(2j)$								
0	1							1
1			1					0
2			1		1			1
3	1		1		1		1	4
4	1		2		1	1	1	5
5	1		2		2	1	2	8
6	1		3	1	3	1	2	9
7	1		4	1	3	2	3	12
8	2		4	1	4	3	4	17
9	2		5	2	5	3	5	20
10	2	1	6	2	6	3	6	24
$\nu_0(f)$	12	1	0	0	26	14	48	101

(81)

4.7 Harmonic analysis on $\mathcal{S}_0(3)$.

We summarize the basis construction for the harmonic analysis on $\mathcal{S}_0(3)$ in terms of C_5 -periodic states on the sphere S^3 .

(i) The spherical harmonics for fixed degree $2j = 0, 1, 2, \dots$ are the Wigner $D^j(u)$ -functions given in eq. 85.

(ii) For the reduction $O(4, R) > S(5)$, the multiplicity of representations D^f is known

from **Table 4.9** and computable from eq. 61 with the characters $\chi^{(j,j)}$ for $O(4, R)$ from **Tables 4.7, 4.8** and χ^f for $S(5)$ given in **Table 4.1**. The partitions $f = [41], [2111]$ are forbidden. For allowed partitions f , the explicit state construction can employ the general Young operators from representation theory, [13] pp. 112-23, with the properties

$$\begin{aligned} c_{r,s}^f &:= \frac{|f|}{5!} \sum_{p' \in S(5)} D_{r,s}^f(p') T_{p'}, \\ T_p c_{r,s}^f &= \sum_{s'} c_{r',s}^f D_{r',r}^f(p) \\ c_{\tilde{s},\tilde{r}}^{\tilde{f}} c_{r,s}^f &= \delta(\tilde{f}, f) \delta(\tilde{r}, r) c_{\tilde{s},s}^f, \\ (c_{r,s}^f)^\dagger &= c_{s,r}^f \end{aligned} \tag{82}$$

The rows and columns (r, s) of D^f can be labelled by Young diagrams or by Yamanouchi symbols as in [3], compare for example eq. 29. The second line in eq. 82 shows that the application of a Young operator to a state yields a basis function for the irreducible representation D^f with row label r . To apply the Young operator to a spherical harmonic, one needs the matrix elements of all permutations $p \in S(5)$ for the representation $D^{(j,j)}$ of $O(4, R)$. Any element $p \in S(5)$ can be written as a product of the generators given in **Table 4.2**. The corresponding product of operators T_p is carried out with the help of eq. 60. Then the expressions eq. 44 or 54 yield the matrix elements $D^{(j,j)}(p)$. The matrix elements in the Young orthogonal representation $D^f(p)$ are computed in a similar way from those of the generators, given for example in [3] pp. 226-30, or constructed directly from [3] pp. 219-20.

(iii) The reduction $S(5) > C_5$ for the 1-dimensional representations $f = [5], [11111]$ is unique, and for the partitions $f = [32], [221], [311]$ by eq. 65 yields one or two unique states with the coefficients eqs. 73, 75, 78 respectively.

The C_5 -periodic basis states on S^3 each belong to a fixed degree $(2j)$ from $O(4, R)$, carry an allowed partition f from $S(5)$, are orthogonal with respect to $(2j)$ and f , and stable under C_5 .

5 Conclusion.

Methods of group theory allow to construct and analyze the harmonic analysis on topological manifolds. This is demonstrated in section 4 for the simplicial manifold $\mathcal{S}_0(3)$. The multiplicities provide the specific selection rules for the chosen simplex topology. The symmetric group $S(5)$ plays a key role. Its representations $f = [41], [2111]$ are eliminated from the harmonic analysis. The details for the basis construction are given in subsection 4.7.

In general, the harmonic analysis on two different manifolds $\mathcal{M}, \mathcal{M}'$ covered by the sphere $S^{(n-1)}$ is unified by the spherical harmonics and corresponding representations.

The differences between topologies appear in the form of different subgroups of deck transformations. In the harmonic analysis these involve different group/subgroup representations and reductions in $O(n, R) > \text{deck}(\mathcal{M})$, $O(n, R) > \text{deck}(\mathcal{M}')$. Intermediate subgroups as $S(n+1)$ in eq. 2 can dominate the harmonic analysis on spherical manifolds. The reduction $O(n, R) > S(n+1)$, $n > 2$ for simplicial manifolds may require generalized Casimir operators as exemplified in [5] for $n = 3$. Selection rules for $S(n+1) > C_{n+1}$ eliminate complete representations D^f of the group $S(n+1)$ from the harmonic analysis on the sphere $S^{(n-1)}$ when restricted to the simplicial manifold.

To see the topological variety of the harmonic analysis, compare the tetrahedral Platonic 3-manifold $\mathcal{M}$ analyzed here with the dodecahedral Platonic 3-manifold $\mathcal{M}'$. The homotopy group of Poincare's dodecahedral 3-manifold $\mathcal{M}'$ is, compare [11] pp. 216-8, the binary icosahedral group. It was found in [6] that the isomorphic group $\text{deck}(\mathcal{M}')$ acts exclusively as a subgroup of $SU(2, C)^r$ from the right on the sphere S^3 in the coordinates eq. 41, with the consequence of a degeneracy of the dodecahedral eigenmodes. The multiplicity in the reduction from $O(4, R)$ to the subset of eigenmodes for the dodecahedral 3-manifold is completely resolved by a generalized Casimir operator. The multiplicity analysis in [6], [7] shows that the lowest dodecahedral eigenmodes are of degree $(2j) = 12$.

Comparison with the present harmonic analysis for the simplicial 3-manifold demonstrates a genuine dependence of the selection rules and the spectrum of eigenmodes on the topology and on the topologically invariant subgroups involved. Corresponding implications can be drawn for the use of harmonic analysis in the cosmic topology of 3-space.

Appendix: Some properties of the group $SU(2, C)$ and its representations.

The group elements $u \in SU(2, C)$ have the unitary, unimodular and complex conjugation properties

$$\begin{aligned} u^\dagger &= u^{-1}, \det(u) = 1, \\ \bar{u} &= q^{-1}uq, \quad q^{-1} = -q = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}, \quad q^T = q^{-1} \end{aligned} \tag{83}$$

The irreducible representations of $SU(2, C)$ are the Wigner D^j functions

$$D_{m_1 m_2}^j(u), j = 0, 1/2, 1, \dots, -j \leq m_i \leq j, i = 1, 2. \tag{84}$$

which are homogeneous polynomials with real coefficients of degree $2j$ in the complex matrix elements eq. 41 of u . They are explicitly given, [13] pp. 163-6, [1] pp. 56-67,

by

$$\begin{aligned}
D_{m_1 m_2}^j(z_1, z_2, \bar{z}_1, \bar{z}_2) &= \left[\frac{(j+m_1)!(j-m_1)!}{(j+m_2)!(j-m_2)!} \right]^{1/2} \\
&\times \sum_{\sigma} \frac{(j+m_2)!(j-m_2)!(-1)^{m_2-m_1+\sigma}}{(j+m_1-\sigma)!(m_2-m_1+\sigma)!\sigma!(j-m_2-\sigma)!} \\
&\times z_1^{j+m_1-\sigma} \bar{z}_2^{m_2-m_1+\sigma} z_2^{\sigma} \bar{z}_1^{j-m_2-\sigma}, \\
j &= 0, 1/2, 1, 3/2, \dots
\end{aligned} \tag{85}$$

Unitarity and reality relations of these polynomial representations imply

$$\begin{aligned}
D_{m_1 m_2}^j(u^{-1}) &= \overline{D_{m_2 m_1}^j(u)}, \\
\overline{D_{m_1 m_2}^j(u)} &= D_{m_1 m_2}^j(\bar{u}).
\end{aligned} \tag{86}$$

From these equations follows the transposition property

$$D_{m_1 m_2}^j(u^T) = D_{m_2 m_1}^j(u). \tag{87}$$

The polynomial homogeneity from eq. 85 reads

$$D_{m_1 m_2}^j(\lambda u) = \lambda^{(2j)} D_{m_1 m_2}^j(u). \tag{88}$$

To pass from an element $g \in SU(2, C)$ to its character $\chi^j(g)$ in the representation $D^j(g)$ we first obtain the angle ϕ from its trace,

$$\frac{1}{2} \chi^{1/2}(g) = \frac{1}{2} \text{Trace}(g) := \cos(\phi/2), \tag{89}$$

and then for general j use

$$\chi^j(g) = \sum_{m=-j}^j \exp(im\phi) = \frac{\sin(\frac{\phi(2j+1)}{2})}{\sin(\frac{\phi}{2})}. \tag{90}$$

References

- [1] Edmonds A R, *Angular momentum in quantum mechanics*, Princeton University Press, Princeton 1957
- [2] Everitt B, *3-manifolds from Platonic solids*, Topology and its Applications **138** (2004), 253-63
- [3] Hamermesh M, *Group theory and its application to physical problems*, Addison-Wesley, Reading 1962

- [4] Humphreys J E, *Reflection groups and Coxeter groups*, Cambridge University Press, Cambridge 1990
- [5] Kramer P and Moshinsky M, *Group Theory of Harmonic Oscillators, III. States with Permutational Symmetry*, Nuclear Physics **82** (1966) 241-74
- [6] Kramer P, *An invariant operator due to F Klein quantizes H Poincare's dodecahedral manifold*, J Phys A: Math Gen **38** (2005) 3517-40
- [7] Kramer P, *Harmonic polynomials on the Poincare dodecahedral 3-manifold* J. of Geometry and Symmetry in Physics **6** (2006) 55-66
- [8] Lachize-Rey M and Luminet J-P, *Cosmic topology*, Phys Rep **254** (1995) 135-214
- [9] Luminet J-P, Weeks J R, Riazuelo A, Lehoucq R and Uzan J-Ph, *Dodecahedral space topology as an explanation for weak wide-angle temperature correlations in the cosmic microwave background*, Nature **425** (2003) 593-5
- [10] Robinson G de B, *Representation theory of the symmetric group*, University of Toronto Press 1961
- [11] Seifert H and Threlfall W, *Lehrbuch der Topologie*, Leipzig 1934, Chelsea Reprint 1980
- [12] Thurston W P, *Three-Dimensional Geometry and Topology*, Princeton University Press, Princeton 1997
- [13] Wigner E P, *Group theory and its application to the quantum mechanics of atomic spectra*, Academic Press, New York (1959)

6 Acknowledgement.

My thanks are due to Marcos Moshinsky, UNAM Mexico, who over many years allowed me to share his insight into groups and their representation theory. Moreover during a visit to Mexico in 2008 he participated in the set-up of the present research project. Many thanks are due to R. Kellerhals, U. Fribourg, Switzerland, for pointing out reference [2], and to T. Kramer, U. Regensburg, Germany, for doing most of the algebraic computations for the Tables.